

EXERCISE SHEET 2

Exercise 1. Let (X, d) be a metric space and $f: X \rightarrow \mathbb{R}$ (note that we do not assume any regularity on f).

- i) For $\epsilon > 0$ let C_ϵ be the set of all $x \in X$ for which there exists a $\delta > 0$ such that, for all $y, z \in X$ with $d(x, y), d(x, z) < \delta$, $|f(y) - f(z)| < \epsilon$. Show that C_ϵ is open.
- ii) Show that the set of points where f is continuous is a countable intersection of open sets.
- iii) Suppose that $X = \mathbb{R}$. Show that the set of points where f is differentiable is a Borel set.

Exercise 2. A subset S of a metric space X is porous if for every $x \in S$ there exists a $\lambda > 0$ and a sequence $x_n \rightarrow x$ such that

$$B(x_n, \lambda d(x, x_n)) \cap S = \emptyset$$

for every $n \in \mathbb{N}$. (Recall that this notion was used in the proof of Rademacher's theorem.)

Prove that any porous subset of a doubling metric measure space has measure zero.

Recall that for a metric space X , $S \subset X$ and $\alpha \geq 0$, the α -dimensional outer Hausdorff measure of S is defined by

$$\mathcal{H}^\alpha(S) = \liminf_{\delta \rightarrow 0} \left\{ \sum_{i \in \mathbb{N}} (\text{diam } S_i)^\alpha : S \subset \bigcup_{i \in \mathbb{N}} S_i, \text{diam } S_i < \delta \right\}.$$

For any metric space X and any $\alpha \geq 0$, \mathcal{H}^α is a Borel measure.

Exercise 3. Let X, Y be metric spaces, $f: X \rightarrow Y$ and $\alpha, L \geq 0$.

- i) Suppose that f is L -Lipschitz and $S \subset X$. Prove that $\mathcal{H}^\alpha(f(S)) \leq L^\alpha \mathcal{H}^\alpha(S)$.
- ii) For $R > 0$ suppose that, for each $x \in S$ and $y \in X$ with $d(x, y) < R$, $d(f(x), f(y)) < Ld(x, y)$. Prove that $\mathcal{H}^\alpha(f(S)) \leq L^\alpha \mathcal{H}^\alpha(S)$.
- iii) Define

$$\text{Lip}(f, x) = \limsup_{y \rightarrow x} \frac{d(f(x), f(y))}{d(x, y)},$$

the pointwise Lipschitz constant of f at x . Suppose that $\text{Lip}(f, x) < L$ for each $x \in X$. Prove that $\mathcal{H}^\alpha(f(X)) \leq L^\alpha \mathcal{H}^\alpha(X)$. (You may assume that $x \mapsto \text{Lip}(f, x)$ is Borel, along with other similar statements.)

Exercise 4. Let $\gamma: [a, b] \rightarrow X$ be Lipschitz.

- i) Prove that $\text{len}(\gamma) \geq d(\gamma(b), \gamma(a))$.
- ii) Prove that the arc-length parametrisation γ^* of any Lipschitz $\gamma: [a, b] \rightarrow X$ exists and is 1-Lipschitz.
- iii) Prove that, for any $s \leq t \in [0, \text{len}(\gamma)]$, $\text{len}(\gamma^*|_{[s, t]}) = t - s$.
- iv) From now on, suppose that γ is injective. Prove that $\text{len}(\gamma^*|_{[s, t]}) = \mathcal{H}^1(\gamma^*([s, t]))$.
- v) Deduce that, for any Borel $S \subset [0, \text{len}(\gamma)]$, $\mathcal{H}^1(\gamma^*(S)) = \mathcal{L}^1(S)$.
- vi) Show that this may be false if γ is not injective.

Exercise 5. Let X, Y be metric spaces, $S \subset X$ and let $f: S \rightarrow Y$ be L -Lipschitz. Suppose that Y is complete. Prove that there exists a unique L -Lipschitz extension of f defined on the closure of S . Prove that this is not necessarily the case if Y is not complete.

Exercise 6. Let X, Y be metric spaces and for each $n \in \mathbb{N}$, $f_n: X \rightarrow Y$ L -Lipschitz. Suppose that $f_n \rightarrow f$ pointwise. Prove that f is L -Lipschitz.

Exercise 7. The space ℓ_∞ consists of all bounded real sequences equipped with the supremum norm. It is a Banach space (that is, a complete vector space; if you have never proved that it is complete, you should do so now).

Let (X, d) be a separable metric space, x_0, x_1, x_2, \dots a dense sequence and define $\iota: X \rightarrow \ell_\infty$ by

$$\iota(x) = (d(x, x_1) - d(x_1, x_0), d(x, x_2) - d(x_2, x_0), d(x, x_3) - d(x_3, x_0), \dots).$$

Prove that ι is well defined and an isometry.

This shows that any separable metric space can be isometrically embedded into a Banach space. However, this Banach space is huge (it is not separable). We can do a little better. Let \tilde{X} be the closed linear span of $\iota(X)$. That is, \tilde{X} is the closure of the set

$$\left\{ \sum_{i=1}^n \lambda_i y_i : n \in \mathbb{N}, \lambda_i \in \mathbb{R}, y_i \in \iota(X) \right\}.$$

Certainly \tilde{X} is a closed subspace (and so a Banach space). Prove that it is separable.